

Original Article

## Home Range Characteristics and Habitat Selection by Daurian Hedgehogs (*Mesechinus dauuricus*) in Ikh Nart Nature Reserve, Mongolia

Mirka Zapletal<sup>1</sup>, Batdorj Sodnompil<sup>2</sup>, Jon Atwood<sup>3</sup>, James D. Murdoch<sup>4</sup> and Richard P. Reading<sup>5</sup>

<sup>1</sup>Antioch University New England, P.O. Box 205, Lincoln, NH 03251, Fax: +1-603-745-2250, Tel.: +1-603-715-4688, e-mail: mirkazapletal@yahoo.com

<sup>2</sup>Mongolian Academy of Sciences, Institute of Biology, Ulaanbaatar 210351, Mongolia, e-mail: Batdorj100@gmail.com

<sup>3</sup>Biodiversity Research Institute, 652 Main Street, Gorham, Maine 04038 USA, e-mail: jon.atwood@briloon.org

<sup>4</sup>University of Vermont, Wildlife and Fisheries Biology Program, Rubenstein School of Environment and Natural Resources, George Aiken Center, Burlington, Vermont 05405 USA, e-mail: jmurdoch@uvm.edu

<sup>5</sup>Denver Zoological Foundation, Conservation Biology Department, 2300 Steele Street, Denver, Colorado 80205 USA, e-mail: rreading@denverzoo.org

### Abstract

---

**Key words:**

compositional analysis, Daurian hedgehog, habitat selection, home range, *Mesechinus*, Mongolia, steppe

**Article information:**

Received: 31 Jul. 2012

Accepted: 04 Dec. 2012

Published: 25 Dec. 2012

**Correspondence:**

mirkazapletal@yahoo.com

We examined home range characteristics and habitat selection of Daurian hedgehogs in Ikh Nart Nature Reserve, Mongolia. Home ranges of hedgehogs varied from 113.15 ha to 2,171.97 ha, and were larger in early summer than late summer. Hedgehogs showed relative preference for rocky outcrops and low-density shrub habitats, and relative avoidance of high-density shrub areas. Habitat selection also changed between early and late summer, shifting to greater use of low-density shrub areas and decreased use of forb-dominated short grass. Our baseline data on home ranges and habitat selection expand understanding of hedgehog ecology and provide guidance for future management decisions in Ikh Nart Nature Reserve and elsewhere in Mongolia.

**Cite this paper as:**

Zapletal, M., Batdorj, S., Atwood, J., Murdoch, J. D. & Reading, R. P., 2012. Home range characteristics and habitat selection by Daurian hedgehogs (*Mesechinus dauuricus*) in Ikh Nart Nature Reserve, Mongolia. *Mong. J. Biol. Sci.*, 10(1-2): 41-50.

---

### Introduction

Daurian hedgehogs (*Mesechinus dauuricus*) are insectivores found in eastern Mongolia and adjacent areas of Siberia and China (Stubbe *et al.*, 2008). Researchers have studied several aspects of European hedgehog (*Erinaceus europaeus*) ecology (Reeve, 1982; Boitani & Reggiani, 1984; Micol *et al.*, 1994; Doncaster *et al.*, 2001; Riber, 2006), but less ecological

research has focused on Daurian hedgehogs. All hedgehogs share a number of anatomical features, such as spines, the orbicularis muscle, and a body-plan described as 'basic' (Reeve, 1994), but behaviors and ecological requirements of populations vary among species and environmental conditions (e.g. food availability, predator density, human impact on the landscape,